

Reproducing Kernel Hilbert Space Technique for Fourth-Order Singular Boundary Value Problems

Khaled K. Jaber

Department of Mathematics, Faculty of Science, Isra University, Amman, Jordan

Email: khaled.jaber@iu.edu.jo

How to cite this paper: Jaber, K.K. (2026) Reproducing Kernel Hilbert Space Technique for Fourth-Order Singular Boundary Value Problems. *Applied Mathematics*, 17, 354-369.

<https://doi.org/10.4236/am.2026.176022>

Received: April 7, 2026

Accepted: June 21, 2026

Published: June 24, 2026

Copyright © 2026 by author(s) and

Scientific Research Publishing Inc.

This work is licensed under the Creative

Commons Attribution International

License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Reproducing kernel Hilbert space method is utilized in this paper as an efficient approach to solve singular fourth order boundary value problems of mixed form Fredholm-Volterra integro-differential equations. In order to obtain the required nodal values, the algorithm developed using two smooth reproducing kernel functions. The solution philosophy is based on applying the Gram-Schmidt process to the kernel function obtained in the space $W_2^5[0,1]$ to produce an orthogonal basis. From that point onward, the orthogonal basis was built for the purpose of formulating and utilizing numerical solutions in the same space. Some linear and nonlinear numerical issues were broken down to delineate the technique and affirm the execution of the proposed strategy. The numerical outcomes emphasize the method role in improving the initial approximation, handling boundary conditions, and refining the solution throughout the iterative process to shed light on such singular equations.

Keywords

Hilbert Space, Integro-Differential Equation, Gram-Schmidt Process, Fredholm-Volterra Integral Equations, Reproducing Kernel Function

1. Introduction

Numerous problems in various fields of science can be diminished, by applying some proper discretization into integro-differential equations (IDEs). In fact, numerous scientific details of engineering applications, power generators, physical phenomena such as fluid mechanics, biological and chemical models contain IDEs with boundary conditions (BCs) etc. [1]-[5]. The term “singular” is used to emphasize that the coefficients of the considered equation may take value zero at one

or more point in the given domain. Anyhow, traditional numerical methods such as finite difference, homotopy analysis and shooting technique encountered loss of accuracy and significant difficulties or may lose the convergence when they applied to problem with singularities. Since it is normally difficult to get the closed frame.

The technique of reproducing kernel has been employed to solve differential equations since the mid-20th century as a novel solver for the BVPs to provide solutions in a structured and systematic manner. In 1907, Zaremba [6] was the first to introduce the kernel corresponding to a class of functions without giving any name or developing any theory, and he was also the first to mention its reproducibility property. In 1909, Mercer [7] developed the theory of continuous positive definite kernels under the name of the theory of “positive definite kernels”. He studied the relationship between positive and negative type functions that satisfy reproducibility property with the theory of integral equations. Moreover, he demonstrated some qualitative characteristics of these positive definite kernels that were not present in all other continuous kernels of IEs.

The goal of numerical techniques is to provide precise strategies to deal with problem in a numerical frame. This technique relies on the use of high-precision computers to perform the steps in order to obtain the result from the raw data.

The RKHS method is a numerical technique developed to solve substantial assortment of ordinary and partial differential equations. The general theory of RKs was systematized in 1950 by Aronszajn [8]. He summarized the previous related research results and used “RK functions” as the identical term for these different functions, so the foundations of RK theory were set up. The applications of the RKHS method in solving different categories of equations with some characteristics and advantages can be found in [9]-[21]. Advances in computer hardware and software make verification and simulation easier, so they are necessary in applied sciences. However, the RKHS method was intentionally examined to solve numerically for the following fourth-order singular BVPs of the mixed-type Fredholm-Volterra IDE in the unknown function u :

$$p_4(x)u^{(4)} + p_3(x)u^{(3)} + p_2(x)u^{(2)} + p_1(x)u^{(1)} = f(x, u) + T[u](x) \quad (1)$$

In which $T[u](x)$ operator is given as

$$T[u](x) = \lambda_1 \int_a^b k_1(x, t) J_1(u(t)) dt + \int_a^x k_2(x, t) J_2(u(t)) dt \quad (2)$$

With respect to the following set of constraint BCs:

$$\begin{aligned} u(a) = \alpha_1, u(b) = \beta_2 \text{ and } u'(a) = \alpha_2, u'(b) = \beta_2 \\ u(a) = \alpha_1, u(b) = \beta_2 \text{ and } u''(a) = \alpha_2, u''(b) = \beta_2 \end{aligned} \quad (3)$$

where $x \in [a, b]$, the function $f(x, u)$ is a continuous real valued function with two variables, λ_1, λ_2 are constant parameters, $k_1(x, t), k_2(x, t)$ are two known kernel functions of two variables x and t , $J_1(u(t)), J_2(u(t))$ are two known functions of one variable, and $T[u](x)$ is Fredholm-Volterra mixed type

operator.

Discussion about solvability of singular BVPs of Fredholm-Volterra mixed type operator subject to different separated BCs is scarce and missing. Lately, some researchers have investigated the numerical applicability of IDEs in the case of nonsingular coefficients by using some of the well-known methods. These methods and their properties can be found in [22]-[26].

Most physical and engineering problems are often described in terms of differential equations, integral equations, or IDEs with conditions imposed at one point or more points see A.Majid Wazwaz book [27].

BVPs of singular functions form emerge as often as possible in numerous branches of science and engineering, such as fluid mechanics, atomic calculations, biological systems, chemical models and in the study of nonlinear elliptic equations see references [28] [29].

In this paper, we present a novel application of the reproducing kernel Hilbert space (RKHS) method to solve linear and nonlinear fourth-order singular BVPs of the mixed-type Fredholm-Volterra integro-differential equations.

2. Preliminaries

Definition 2.1. The inner product and the norm of the space $W_2^1[0,1] = \{z(x) : z \text{ is real-valued function which is absolutely continuous on } [0,1] \text{ and } z' \in L^2[0,1]\}$ are defined respectively as

$$\langle z_1(x), z_2(x) \rangle_{W_2^1} = z_1(0)z_2(0) + \int_0^1 z_1'(x)z_2'(x)dx$$

$$\|z\|_{W_2^1} = \sqrt{\langle z(x), z(x) \rangle_{W_2^1}}$$

where $z_1, z_2 \in W_2^1[0,1]$.

Theorem 2.1. The space $W_2^1[0,1]$ is a RKHS. Using Mathematica software, it was found that its reproduction kernel function $R_x(y)$ is

$$R_x(y) = \begin{cases} R_{1x}(y) = 1+y, & y \leq x, \\ R_{2x}(y) = 1+x, & y > x, \end{cases}$$

Definition 2.2. The inner product and the norm of the space $W_2^5[0,1] = \{z(x) : z^{(i)}, i = 0, 1, 2, 3, 4\} \text{ are real-valued functions that are absolutely continuous on } [0,1], z^{(5)} \in L^2[0,1], \text{ and } z(0) = z'(0) = z(1) = z'(1) = 0\}$ are defined respectively as

$$\langle z_1(x), z_2(x) \rangle_{W_2^5} = \sum_{i=0}^2 z_1^{(i)}(0)z_2^{(i)}(0) + \sum_{i=0}^1 z_1^{(i)}(1)z_2^{(i)}(1) + \int_0^1 z_1^{(5)}(x)z_2^{(5)}(x)dx$$

$$\|z\|_{W_2^5} = \sqrt{\langle z(x), z(x) \rangle_{W_2^5}}$$

where $z_1, z_2 \in W_2^5[0,1]$.

Theorem 2.2. The space $W_2^5[0,1]$ is a RKHS. Using Mathematica, it was found that its reproduction kernel function $K_x(y)$ is

$$K_x(y) = \begin{cases} K_{1x}(y) = \sum_{i=0}^9 p_i(x) y^i, & y \leq x, \\ K_{2x}(y) = \sum_{i=0}^9 q_i(x) y^i, & y > x, \end{cases}$$

where

$$\begin{aligned} K_{1x}(y) &= \frac{1}{362880} (x-1)^2 y^2 \times (5x^3(8-15y)y + x^6(19-12y)y + y^7 \\ &\quad + xy^6(-9+2y) + 3x^5y(-10+3y) + x^7y(-4+3y) + 5x^4y(1+6y) \\ &\quad + 3x^2(30240-60480y+30240y^2+12y^5-6y^6+y^7)) \\ K_{2x}(y) &= \frac{1}{362880} x^2 (y-1)^2 \times (90720y^2 + 36x^5y^2 - 9x^6y(1+2y) \\ &\quad + x^7(1+2y+3y^2) + xy^2(-181440+40y+5y^2-30y^3+19y^4-4y^5) \\ &\quad + 3x^2y^2(30240-25y+10y^2+3y^3-4y^4+y^5)) \end{aligned}$$

3. Problem Formulation

The key aspect of the process is selecting appropriate linear operator depending on the inner product spaces $W_2^5[0,1]$ and $W_2^1[0,1]$, this part is the problem formulating. The nonhomogenous BC's must be converted using appropriate transformation to homogeneous BC's in order to apply the RKHS method. Denote the solution of the new equation by $u(x)$. we get

$$p_4(x)u^{(4)} + p_3(x)u^{(3)} + p_2(x)u^{(2)} + p_1(x)u^{(1)} = f(x, u) + T[u](x)$$

where $T[u](x)$ is given by

$$T[u](x) = \lambda_1 \int_0^1 k_1(x, t) J_1(u(t)) dt + \lambda_2 \int_0^x k_2(x, t) J_2(u(t)) dt$$

subject to the following constraint boundary conditions:

$$\begin{aligned} u(0) &= u'(0) = 0, \\ u(1) &= u'(1) = 0. \end{aligned}$$

To perform the procedure, we introduce the operator

$$L : W_2^5[0,1] \rightarrow W_2^1[0,1] \quad (4)$$

as

$$Lu(x) = p_4(x)u^{(4)} + p_3(x)u^{(3)} + p_2(x)u^{(2)} + p_1(x)u^{(1)} \quad (5)$$

Thus, discretized form of equivalent equations can be obtained as follows:

$$Lu(x) = F(x, u(x), T[u](x)) \quad (6)$$

subject to the boundary conditions

$$\begin{aligned} u(0) &= u'(0) = 0, \\ u(1) &= u'(1) = 0. \end{aligned} \quad (7)$$

in which the function F is constructed as follows

$$F(x, u(x), T[u](x)) = f(x, u) + T[u](x) \quad (8)$$

Theorem 3.1. The operator $L : W_2^5[0, 1] \rightarrow W_2^1[0, 1]$ is bounded and linear.

Proof. The lineart part is obvious. From definition 2.2. we have

$$\|(Lu)(x)\|_{W_2^1}^2 = [(Lu)(0)]^2 + \int_0^1 [(Lu)'(x)]^2 dx \quad (9)$$

Using the reproducing kernel function properties we can write

$$\begin{aligned} v(x) &= \langle v(y), K_x(y) \rangle_{W_2^5} \\ (Lv)(x) &= \langle v(y), (LK_x)(y) \rangle_{W_2^5} \\ (Lv)'(x) &= \langle v(y), (LK_x)'(y) \rangle_{W_2^5} \end{aligned}$$

Schwarz inequality gives

$$\begin{aligned} |(Lu)(x)| &= \left| \langle u(x), (LK_x)(x) \rangle_{W_2^5} \right| \\ &\leq \|LK_x(x)\|_{W_2^5} \|u(x)\|_{W_2^5} \\ &= M_1 \|u(x)\|_{W_2^5} \\ |(Lu)'(x)| &= \left| \langle v(x), (LK_x)'(x) \rangle_{W_2^5} \right| \\ &\leq \|(LK_x)'(x)\|_{W_2^5} \|u(x)\|_{W_2^5} \\ &= M_2 \|u(x)\|_{W_2^5} \end{aligned}$$

Thus, after simple computations, we get

$$\begin{aligned} \|(Lu)(x)\|_{W_2^1}^2 &= [(Lu)(0)]^2 + \int_0^1 [(Lu)'(x)]^2 dx \\ &\leq (M_1^2 + M_2^2) \|u(x)\|_{W_2^5}^2 \end{aligned}$$

or $\|(Lv)(x)\|_{W_2^1} \leq M \|v(x)\|_{W_2^5}$, where $M = \sqrt{M_1^2 + M_2^2}$.

Theorem 3.2. The sequense $\{\psi_i(x)\}_{i=1}^\infty$ is a complete function system for the space $W_2^5[0, 1]$ and

$$\psi_i(x) = L_y K_x(y) \Big|_{y=x_i}.$$

Proof.

$$\begin{aligned} \psi_i(x) &= L_i^* \varphi(x) \\ &= \langle L_i^* \varphi(y), K_x(y) \rangle_{W_2^5} \\ &= \langle \varphi_i(y), L_y K_x(y) \rangle_{W_2^1} \\ &= L_y K_x(y) \Big|_{y=x_i}. \end{aligned}$$

For each $u(x) \in W_2^5[0, 1]$, let $\langle u(x), \psi_i(x) \rangle_{W_2^5} = 0$, $\forall i \in \mathbb{N}$. Then,

$$\langle u(x), \psi_i(x) \rangle_{W_2^5} = \langle u(x), L^* \varphi_i(x) \rangle_{W_2^5} = \langle Lu(x), \varphi_i(x) \rangle_{W_2^1} = Lu(x_i) = 0 \quad (10)$$

Therefore $Lu(x) = 0$. Since L has a trivial kernel under the given boundary conditions, $u(x) = 0$. Using Gram-Schmidt orthogonalization process the series $\{\bar{\psi}_i(x)\}_{i=1}^{\infty}$ is given by

$$\bar{\psi}_i(x) = \sum_{k=1}^i \beta_{ik} \psi_k(x) \quad (11)$$

where

$$\beta_{ij} = \begin{cases} \frac{1}{\|\psi_1\|_{W_2^5}} & i = j = 1 \\ \frac{1}{d_{ik}} & i = j \neq 1 \\ -\frac{1}{d_{ik}} \sum_{k=j}^{i-1} c_{ik} \beta_{kj} & i > j \end{cases}$$

where $d_{ik} = \sqrt{\|\psi_i\|_{W_2^5}^2 - \sum_{k=1}^{i-1} c_{ik}^2}$ and $c_{ik} = \langle \psi_i, \bar{\psi}_k \rangle_{W_2^5}$.

Lemma 3.1. For each $i = 0, 1, 2, 3, 4$ there exist a positive M_i such that $\|u^{(i)}(x)\|_C \leq M_i \|u(x)\|_{W_2^5}$, where

$$\|u(x)\|_C = \max_{0 \leq x \leq 1} |u(x)|$$

Proof. $\forall x \in [0, 1]$, we have

$$u^{(i)}(x) = \langle u(y), K_x^{(i)}(y) \rangle_{W_2^5}, i = 0, 1, 2, 3, 4$$

By the reproduction kernel function $K_x(y)$ given in theorem 2.2 it follows that for each $i = 0, 1, 2, 3, 4$

$$\|K_x^{(i)}(y)\|_{W_2^5} \leq M_i$$

Thus,

$$\begin{aligned} |u^{(i)}(x)| &= \left| \langle u(x), K_x^{(i)}(x) \rangle_{W_2^5} \right| \\ &\leq \|K_x^{(i)}(x)\|_{W_2^5} \|u(x)\|_{W_2^5} \\ &\leq M_i \|u(x)\|_{W_2^5} \end{aligned}$$

Hence, for each $i = 0, 1, 2, 3, 4$

$$\|u^{(i)}(x)\|_C \leq M_i \|u(x)\|_{W_2^5} \quad (12)$$

Theorem 3.3. The exact solution of Equations (5) and (6) is given by

$$u(x) = \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} F(x_k, u(x_k), Tu(x_k)) \bar{\psi}_i(x)$$

Proof. Let $u(x)$ be solution. Then, using Equation (11), we have

$$\begin{aligned}
u(x) &= \sum_{i=1}^{\infty} \langle u(x), \bar{\psi}_i(x) \rangle_{W_2^5} \bar{\psi}_i(x) \\
&= \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} \langle u(x), \psi_k(x) \rangle_{W_2^5} \bar{\psi}_i(x) \\
&= \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} \langle u(x), L^* \varphi_k(x) \rangle_{W_2^5} \bar{\psi}_i(x) \\
&= \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} \langle Lu(x), \varphi_k(x) \rangle_{W_2^1} \bar{\psi}_i(x) \\
&= \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} \langle F(x, u(x), Tu(x)), \varphi_k(x) \rangle_{W_2^1} \bar{\psi}_i(x)
\end{aligned}$$

So, we have

$$u(x) = \sum_{i=1}^{\infty} \sum_{k=1}^i \beta_{ik} F(x_k, u(x_k), Tu(x_k)) \bar{\psi}_i(x) \quad (13)$$

To approximate the solution given in Equation (13), the first n -terms are taken

$$u_n(x) = \sum_{i=1}^n \sum_{k=1}^i \beta_{ik} F(x_k, u(x_k), Tu(x_k)) \bar{\psi}_i(x) \quad (14)$$

Theorem 3.4. The approximated solution $u_n(x)$ converges uniformly to the exact solution $u(x)$. Also, for each $i = 0, 1, 2, 3, 4$ $u_n^{(i)}(x) \rightarrow u^{(i)}(x)$ uniformly.

Proof. By Lemma 3.1, $\forall x \in [a, b]$

$$\begin{aligned}
|u_n(x) - u(x)| &= \left| \langle u_n(x) - u(x), K_x(x) \rangle_{W_2^5} \right| \\
&\leq \|K_x(x)\|_{W_2^5} \|u_n(x) - u(x)\|_{W_2^5} \\
&\leq M_0 \|u_n(x) - u(x)\|_{W_2^5}
\end{aligned}$$

For the derivatives,

$$\begin{aligned}
|u_n^{(i)}(x) - u^{(i)}(x)| &= \left| \langle u_n(x) - u(x), K_x^{(i)}(x) \rangle_{W_2^5} \right| \\
&\leq \|K_x^{(i)}(x)\|_{W_2^5} \|u_n(x) - u(x)\|_{W_2^5} \\
&\leq M_i \|u_n(x) - u(x)\|_{W_2^5}, \quad M_i > 0, \quad i = 1, 2, 3, 4
\end{aligned}$$

Hence, if $\|u_n(x) - u(x)\|_{W_2^5} \rightarrow 0$ as $n \rightarrow \infty$, then $u_n(x) \rightarrow u(x)$ and $u_n^{(i)}(x) \rightarrow u^{(i)}(x)$ uniformly, $i = 1, 2, 3, 4$.

4. Representation of Numerical Solution

In this section, we are giving the numerical solutions form of Equations (5) and (6) in the space $W_2^5[0, 1]$. Then, an iterative formulas for approximate solution is presented.

So as to inherit the behaviour of the numerical solution in the presented RKHS method the following results are guarantee this important requirements.

Lemma 4.1. If $u_n(x) \rightarrow u(x)$ in the sense of the norm of $W_2^5[0, 1]$, $x_n \rightarrow y$ as $n \rightarrow \infty$, then as $n \rightarrow \infty$, we have

$$F(x_n, u_{n-1}(x_n), Tu_{n-1}(x_n)) \rightarrow F(y, u(y), Tu(y))$$

Proof. In the first, we will show that $u_{n-1}(x_n) \rightarrow u(y)$ in the sense of the norm defined on the space $W_2^5[0,1]$. Since

$$|u_{n-1}(x_n) - u(y)| = |u_{n-1}(x_n) - u_{n-1}(y) + u_{n-1}(y) - u(y)|$$

By triangle inequality and the reproducing property of $K_x(y)$, we get

$$\begin{aligned} |u_{n-1}(x_n) - u_{n-1}(y)| &= \left| \langle u_{n-1}(x), K_{x_n}(x) - K_y(x) \rangle_{W_2^5} \right| \\ &\leq \|u_{n-1}(x)\|_{W_2^5} \|K_{x_n}(x) - K_y(x)\|_{W_2^5} \end{aligned}$$

So, we conclude that $\|K_{x_n}(x) - K_y(x)\|_{W_2^5} \rightarrow 0$ as $n \rightarrow \infty$.

Hence, $|u_{n-1}(x_n) - u_{n-1}(y)| \rightarrow 0$ as $x_n \rightarrow y$. Thus, for any $y \in [0,1]$, it holds that $|u_{n-1}(y) - u(y)| \rightarrow 0$ as $n \rightarrow \infty$. Therefore, $u_{n-1}(x_n) \rightarrow u(y)$ as $n \rightarrow \infty$.

Thus, the continuity of J_1 , J_2 , and f , it is obtained that

$$\begin{aligned} J_1(u_{n-1}(x_n)) &\rightarrow J_1(u(y)), \\ J_2(u_{n-1}(x_n)) &\rightarrow J_2(u(y)), \\ f(x_n, u_{n-1}(x_n)) &\rightarrow f(y, u(y)) \end{aligned}$$

This shows that $Tu_{n-1}(x_n) \rightarrow Tu(y)$ as $n \rightarrow \infty$. Hence, we get the results.

The following theorem can be proved similarly.

Theorem 4.1. In the space $W_2^5[0,1]$ where $u_n(x)$ and $u(x)$ given in Equation (13) and Equation (14) we have

1) $u_n(x) \rightarrow u(x)$. Since $\{\psi_i\}$ is complete in $W_2^5[0,1]$, the partial sums u_n converge to u in the norm of $W_2^5[0,1]$.

2) Let $\varepsilon_n^2 = \|u(x) - u_n(x)\|_{W_2^5}^2$ for each natural number n . Then, the sequence of numbers $\{\varepsilon_n\}$ is monotonic decreasing and converges to zero in the sense of the norm of $W_2^5[0,1]$.

Numerical Results

We are giving the complete numerical process tend to emphasize the implementation of algorithms.

Algorithm 4.1. To homogenized the constraints boundary conditions in Equation (3), we do the following:

Step 1: define $v(x)$ as

$$v(x) = u(x) - \phi(x)$$

where $\phi(x)$ is a function satisfying the conditions

$$\phi(0) = \alpha_1, \quad \phi(1) = \beta_1, \quad \phi'(0) = \alpha_2 \quad \text{and} \quad \phi'(1) = \beta_2.$$

Step 2: After easy calculations, one can obtain

$$\phi(x) = \left(\beta_1 - \alpha_1 - \alpha_2 - \frac{3\beta_1 - \beta_2}{2\alpha_2 + 3\alpha_1} \right) x^3 + \frac{3\beta_1 - \beta_2}{2\alpha_2 + 3\alpha_1} x^2 + \alpha_2 x + \alpha_1$$

Step 3: Substitute the new presented function $v(x)$ into Equations (1), (2) and boundary conditions (3) the problem transformed to the following IDE with ho-

mogenous BC's.

$$\begin{aligned} & p_4(x)v^{(4)} + p_3(x)(v^{(3)} + \phi^{(3)})(x) + p_2(x)(v^{(2)} + \phi^{(2)})(x) + p_1(x)((v^{(1)} + \phi^{(1)})(x)) \\ & = f(x, (v + \phi)(x)) + T[(v + \phi)](x) \end{aligned}$$

subject to the homogeneous boundary conditions

$$\begin{aligned} v(0) &= v(1) = 0, \\ v''(0) &= v''(1) = 0. \end{aligned}$$

Step 4: Write known function $v(x)$ as

$$u(x) = v(x) + \phi(x)$$

We mention here that this transformation will be applying on the numerical solution as well as the exact solution.

To solve numerically Equations (1) and (2) subject to the boundary conditions (3) using the RKHS method we implement the following procedure.

Algorithm 4.2. To approximate the solution $u_n(t)$ of $u(t)$, we do the following:

Step a: Choose n assembling points in $[0, 1]$;

Step b: Set $\psi_i(t) = L_s R_t(s) \Big|_{s=t_i}$;

Step c: Find the coefficients β_{ik} ;

Step d: Set $\bar{\psi}_i(x) = \sum_{k=1}^i \beta_{ik} \psi_k(x)$, for $i = 1, 2, \dots, n$;

Step e: Set $i = 1$;

Step f: Set $u_i(x) = \sum_{j=1}^i \sum_{k=1}^j \beta_{ik} F(x_k, u(x_k), Tu(x_k)) \bar{\psi}_j(x)$;

Step g: If $i < n$, then set $i = i + 1$ and go to step 6, else stop.

Example 1. Consider the linear equation:

$$x^4 u^{(4)}(x) - xu'''(x) + u''(x) - \sin(x)u'(x) + u(x) = f(x) + T[u](x)$$

where

$$T[u](x) = \int_0^1 (xt - 1)u(t)dt + \int_0^x \cosh(t)x^3 u(t)dt$$

subject to the boundary conditions

$$\begin{aligned} u(0) &= u''(0) = 1, \\ u(1) &= u''(1) = \cosh(1), \end{aligned}$$

where $0 \leq t < x \leq 1$ and $f(x)$ is chosen such that the exact solution is

$$u(x) = \cosh(x)$$

The approximate solution obtained using the RKM4 method is shown in **Figure 1**, while its derivative is illustrated in **Figure 2** and the corresponding numerical solution is tabulated in **Table 1**.

Example 2. Consider the nonlinear equation:

$$x(x-1)u^{(4)}(x) - xu'''(x) - u(x)e^{u(x)} = f(x) + T[u](x)$$

where

$$T[u](x) = \int_0^1 e^x (t-1) \sinh(u(x)) dt + \int_0^x \cosh(x) t^3 e^{u(t)} dt$$

subject to the boundary conditions

$$\begin{aligned} u(0) &= 0, u''(0) = 1, \\ u(1) &= \ln(2), u''(1) = -\frac{1}{4}, \end{aligned}$$

where $0 \leq t < x \leq 1$ and $f(x)$ is chosen such that the exact solution is

$$u(x) = \ln(x+1)$$

The approximate solution obtained using the RKM4 method is shown in **Figure 3**, while its derivative is illustrated in **Figure 4** and the corresponding numerical solution is tabulated in **Table 2**.

Example 3. Consider the following nonlinear equation:

$$\sinh(x)u^{(4)}(x) - xu''(x) - (x-1)u'(x) + u^3(x) = f(x) + T[u](x),$$

in which the mixed operator is given as

$$T[u](x) = \int_0^1 (x-1) \ln(u(t)) dt + \int_0^x (x+1)^2 t u^3(t) dt$$

and subject to the boundary conditions

$$\begin{aligned} u(0) &= 1, u(1) = e, \\ u'(0) &= 1, u'(1) = e, \end{aligned}$$

where $0 \leq t < x \leq 1$ and $f(x)$ is chosen such that the exact solution is

$$u(x) = e^{-x}.$$

The approximate solution obtained using the RKM4 method is shown in **Figure 5**, while its derivative is illustrated in **Figure 6** and the corresponding numerical solution is tabulated in **Table 3**.

Using all the previous Algorithms, taking $x_i = \frac{i-1}{n-1}$, $i = 1, 2, \dots, n$. The approximate solutions $u_n(x)$ of $u(x)$ at some specified points for $n = 101$ are computed and tabulated.

Table 1. Numerical outcomes for Example1.

x	Exact solution	Approximate solution	Absolute error
0	1	1	0
0.1	1.00500417	1.005004305	$1.35377662 \times 10^{-7}$
0.2	1.02006676	1.020074585	$7.82499771 \times 10^{-6}$
0.3	1.04533851	1.045348268	$9.75802745 \times 10^{-6}$
0.4	1.08107237	1.081079438	$7.06834965 \times 10^{-6}$
0.5	1.12762597	1.127635882	$9.91155212 \times 10^{-6}$
0.6	1.18546522	1.185474920	$9.69967018 \times 10^{-6}$
0.7	1.25516901	1.255175274	$6.26447249 \times 10^{-6}$

Continued

0.8	1.33743495	1.337438958	$4.00825596 \times 10^{-6}$
0.9	1.43308639	1.433089170	$2.77962568 \times 10^{-6}$
1	1.54308063	1.543080630	0

Table 2. Numerical outcomes for Example 2.

x	Exact solution	Approximate solution	Absolute error
0	0	0	0
0.1	0.09531018	0.095323680	$1.34997372 \times 10^{-5}$
0.2	0.18232156	0.182337252	$1.56924138 \times 10^{-5}$
0.3	0.26236426	0.262480352	$1.16091509 \times 10^{-4}$
0.4	0.33647224	0.336572943	$1.00703178 \times 10^{-4}$
0.5	0.40546511	0.405647270	$1.82160060 \times 10^{-4}$
0.6	0.47000363	0.470080576	$7.69462790 \times 10^{-5}$
0.7	0.53062825	0.531134512	$5.06262154 \times 10^{-4}$
0.8	0.58778666	0.587902612	$1.15951598 \times 10^{-4}$
0.9	0.64185389	0.642102879	$2.48989019 \times 10^{-4}$
1	0.69314718	0.693147180	0

Table 3. Numerical results for Example 3.

x	Exact solution	Approximate solution	Absolute error
0	1	1	0
0.1	0.90483742	0.904842776	$5.35609058 \times 10^{-6}$
0.2	0.81873075	0.818733318	$2.56787841 \times 10^{-6}$
0.3	0.74081822	0.740819437	$1.21697472 \times 10^{-6}$
0.4	0.67032005	0.670354405	$3.43549428 \times 10^{-5}$
0.5	0.60653066	0.606548251	$1.75913076 \times 10^{-5}$
0.6	0.54881164	0.548875657	$6.40170534 \times 10^{-5}$
0.7	0.49658530	0.496598855	$1.35553766 \times 10^{-5}$
0.8	0.44932896	0.449382572	$5.36120945 \times 10^{-5}$
0.9	0.40656966	0.406636500	$6.68397647 \times 10^{-5}$
1	0.36787944	0.367879440	0

Next, the geometric behaviours of the approximate solutions and the first derivative of the approximate solutions using the RKHS method are described for all listed problems, respectively.




Figure 1. Skeeth of the approximate solution using the RKHS method in Example 1.




Figure 2. Skeeth of the first derivative of the approximate solution using the RKHS method in Example 1.




Figure 3. Skeeth of the approximate solution using the RKHS method in Example 2.




Figure 4. Skech of the first derivative of the apptoximate solution using the RKHS method in Example 2.




Figure 5. Skech of the apptoximate solution using the RKHS method in Example 3.




Figure 6. Skech of the first derivative of the apptoximate solution using the RKHS method in Example 3.

5. Conclusions

This work was proposed and applied the RKHS method to solve three benchmark singular boundary value problems. The solution's methodology is grounded in generating orthonormal basis functions derived from the reproducing kernels. This orthonormal basis is constructed to facilitate the formulation and employ numerical solutions. Also, this constructed basis provides efficiently convergent approximate solutions. Additionally, both the approximate solution and its derivatives uniformly approach the exact solution and its respective derivatives.

The numerical results demonstrate the efficiency, reliability, and validity of the RKHS method, highlighting its strength and ease of handling these test cases. Meanwhile, from the given data, one can conclude that the validity of any numerical method depends on the complexity of the problem discussed.

Conflicts of Interest

The author declares no conflicts of interest regarding the publication of this paper.

References

- [1] Kanwal, R.P. (1997) Linear Integral Equations Theory and Technique.
- [2] Wazwaz, A. (2006) A Comparison Study between the Modified Decomposition Method and the Traditional Methods for Solving Nonlinear Integral Equations. *Applied Mathematics and Computation*, **181**, 1703-1712. <https://doi.org/10.1016/j.amc.2006.03.023>
- [3] Bloom, F. (1980) Asymptotic Bounds for Solutions to a System of Damped Integrodifferential Equations of Electromagnetic Theory. *Journal of Mathematical Analysis and Applications*, **73**, 524-542. [https://doi.org/10.1016/0022-247x\(80\)90297-8](https://doi.org/10.1016/0022-247x(80)90297-8)
- [4] Holm aker, K. (1993) Global Asymptotic Stability for a Stationary Solution of a System of Integro-Differential Equations Describing the Formation of Liver Zones. *SIAM Journal on Mathematical Analysis*, **24**, 116-128. <https://doi.org/10.1137/0524008>
- [5] Doddrell, D.M., Forbes, L.K. and Crozier, S. (1997) Calculating Current Densities and Fields Produced by Shielded Magnetic Resonance Imaging Probes. *SIAM Journal on Applied Mathematics*, **57**, 401-425. <https://doi.org/10.1137/s0036139995283110>
- [6] Zaremba, S. (1907) L'equation biharmonique et une class remarquable defonctions fondamentals harmoniques. *Bulletin International de l'Academie des Sciences de Cracovie*, 147-196. https://www.biodiversitylibrary.org/item/276656?utm_source=chatgpt.com#page/11/mode/1up
- [7] Mercer, J. (1909) XVI. Functions of Positive and Negative Type, and Their Connection the Theory of Integral Equations. *Philosophical Transactions of the Royal Society of London. Series A, Containing Papers of a Mathematical or Physical Character*, **209**, 415-446. <https://doi.org/10.1098/rsta.1909.0016>
- [8] Aronszajn, N. (1950) Theory of Reproducing Kernels. *Transactions of the American Mathematical Society*, **68**, 337-404. <https://doi.org/10.1090/s0002-9947-1950-0051437-7>
- [9] Cui, M. and Lin, Y. (2009) Nonlinear Numerical Analysis in the Reproducing Kernel Space. Nova Science.

- [10] Berline, A. and Agnan, C.T. (2004) Reproducing Kernel Hilbert Space in Probability and Statistics. Kluwer Academic Publishers.
<https://doi.org/10.1007/978-1-4419-9096-9>
- [11] Daniel, A. (2003) Reproducing Kernel Spaces and Applications. Springer.
- [12] Lin, Y.Z., Cui, M.G. and Yang, L.H. (2006) Representation of the Exact Solution for a Kind of Nonlinear Partial Differential Equations. *Applied Mathematics Letters*, **19**, 808-813.
- [13] Wang, W., Han, B. and Yamamoto, M. (2013) Inverse Heat Problem of Determining Time-Dependent Source Parameter in Reproducing Kernel Space. *Nonlinear Analysis: Real World Applications*, **14**, 875-887.
<https://doi.org/10.1016/j.nonrwa.2012.08.009>
- [14] Jiang, W. and Lin, Y. (2010) Approximate Solution of the Fractional Advection-Dispersion Equation. *Computer Physics Communications*, **181**, 557-561.
<https://doi.org/10.1016/j.cpc.2009.11.004>
- [15] Jiang, W. and Chen, Z. (2014) A Collocation Method Based on Reproducing Kernel for a Modified Anomalous Subdiffusion Equation. *Numerical Methods for Partial Differential Equations*, **30**, 289-300. <https://doi.org/10.1002/num.21809>
- [16] Arqub, O.A. (2016) Approximate Solutions of DASS with Nonclassical Boundary Conditions Using Novel Reproducing Kernel Algorithm. *Fundamenta Informaticae*, **146**, 231-254. <https://doi.org/10.3233/fi-2016-1384>
- [17] Jiang, W. and Chen, Z. (2013) Solving a System of Linear Volterra Integral Equations Using the New Reproducing Kernel Method. *Applied Mathematics and Computation*, **219**, 10225-10230. <https://doi.org/10.1016/j.amc.2013.03.123>
- [18] Arqub, O.A., Al-Smadi, M. and Shawagfeh, N. (2013) Solving Fredholm Integro-Differential Equations Using Reproducing Kernel Hilbert Space Method. *Applied Mathematics and Computation*, **219**, 8938-8948.
<https://doi.org/10.1016/j.amc.2013.03.006>
- [19] Abu Arqub, O. and Al-Smadi, M. (2014) Numerical Algorithm for Solving Two-Point, Second-Order Periodic Boundary Value Problems for Mixed Integro-Differential Equations. *Applied Mathematics and Computation*, **243**, 911-922.
<https://doi.org/10.1016/j.amc.2014.06.063>
- [20] Geng, F.Z. and Qian, S.P. (2013) Reproducing Kernel Method for Singularly Perturbed Turning Point Problems Having Twin Boundary Layers. *Applied Mathematics Letters*, **26**, 998-1004. <https://doi.org/10.1016/j.aml.2013.05.006>
- [21] Geng, F.Z., Qian, S.P. and Li, S. (2014) A Numerical Method for Singularly Perturbed Turning Point Problems with an Interior Layer. *Journal of Computational and Applied Mathematics*, **255**, 97-105. <https://doi.org/10.1016/j.cam.2013.04.040>
- [22] Geng, F.Z. and Qian, S.P. (2015) Modified Reproducing Kernel Method for Singularly Perturbed Boundary Value Problems with a Delay. *Applied Mathematical Modelling*, **39**, 5592-5597. <https://doi.org/10.1016/j.apm.2015.01.021>
- [23] Wazwaz, A. (2001) A Reliable Algorithm for Solving Boundary Value Problems for Higher-Order Integro-Differential Equations. *Applied Mathematics and Computation*, **118**, 327-342. [https://doi.org/10.1016/S0096-3003\(99\)00225-8](https://doi.org/10.1016/S0096-3003(99)00225-8)
- [24] Arikoglu, A. and Ozkol, I. (2005) Solution of Boundary Value Problems for Integro-Differential Equations by Using Differential Transform Method. *Applied Mathematics and Computation*, **168**, 1145-1158. <https://doi.org/10.1016/j.amc.2004.10.009>
- [25] Yildirim, A. (2008) Solution of BVPs for Fourth-Order Integro-Differential Equations by Using Homotopy Perturbation Method. *Computers & Mathematics with*

-
- Applications*, **56**, 3175-3180. <https://doi.org/10.1016/j.camwa.2008.07.020>
- [26] Hashim, I. (2006) Adomian Decomposition Method for Solving BVPs for Fourth-Order Integro-Differential Equations. *Journal of Computational and Applied Mathematics*, **193**, 658-664. <https://doi.org/10.1016/j.cam.2005.05.034>
- [27] Wazwaz, A.M. (2001) Linear and Nonlinear Integral Equations Methods and Applications. Springer.
- [28] Çağlar, H., Çağlar, N. and Özer, M. (2009) B-Spline Solution of Non-Linear Singular Boundary Value Problems Arising in Physiology. *Chaos, Solitons & Fractals*, **39**, 1232-1237. <https://doi.org/10.1016/j.chaos.2007.06.007>
- [29] Hiltmann, P. and Lory, P. (1983) On Oxygen Diffusion in a Spherical Cell with Michaelis-Menten Oxygen Uptake Kinetics. *Bulletin of Mathematical Biology*, **45**, 661-664. [https://doi.org/10.1016/s0092-8240\(83\)80019-6](https://doi.org/10.1016/s0092-8240(83)80019-6)