

TABLE OF CONTENTS

Volume 6, Number 3A, March 2013

MRI is a state of art imaging modality in characterization of indeterminate adnexal masses	
A. N. D. Dwivedi, S. Jain, R. C. Shukla, M. Jain, A. Srivastava, A. Verma.....	309
Cross-modal connectivity of the secondary auditory cortex with higher visual area in the congenitally deaf—A case study	
Y.-W. Sung, S. Ogawa.....	314
Lesion contrast differences in MRI sequences in multiple sclerosis: Correlation to clinical disability	
M. Rossi, M. Raunio, P. Ryymin, I. Elovaara, P. Dastidar.....	319
Automated measurement of three-dimensional cerebral cortical thickness in Alzheimer’s patients using localized gradient vector trajectory in fuzzy membership maps	
C. Tokunaga, H. Arimura, T. Yoshiura, T. Ohara, Y. Yamashita, K. Kobayashi, T. Magome, Y. Nakamura, H. Honda, H. Hirata, M. Ohki, F. Toyofuku.....	327
Reliability and sensitivity to change of IW-TSE versus DESS magnetic resonance imaging sequences in the assessment of bone marrow lesions in knee osteoarthritis patients: Longitudinal data from the Osteoarthritis Initiative (OAI) cohort	
J.-P. Raynauld, L. M. Wildi, F. Abram, T. Moser, J.-P. Pelletier, J. Martel-Pelletier.....	337
A diffusion-weighted imaging based diagnostic system for early detection of prostate cancer	
A. Firjani, A. Elnakib, F. Khalifa, G. Gimel’farb, M. A. El-Ghar, A. Elmaghraby, A. El-Baz.....	346
Combining DCE-MRI and 1H-MRS spectroscopy by <i>distribution free</i> approach results in a high performance marker: Initial study in breast patients	
S. Nicolosi, G. Russo, I. D’Angelo, G. Vicari, M. C. Gilardi, G. Borasi.....	357
Significance and utility of MR cisternography for diagnosis and treatment of trigeminal neuralgia	
T. Tanaka, S. Shiiba, E. Sakamoto, M. Oda, S. Kito, K. Ono, N. Wakasugi-Sato, S. Matsumoto-Takeda, Y. Seta, Y. Imamura, S. Watanabe, K. Inenaga, Y. Morimoto.....	365
Normalized intervertebral disc MRI signal as a biomarker of pain	
M. Aissiou, D. P éri é J.-M. Mac-Thiong.....	372
Probabilistic tractography of the posterior subthalamic area in Parkinson’s disease patients	
M. G. Garc ía-Gomar, L. Concha, S. Alcauter, J. S. Abraham, J. D. Carrillo-Ruiz, G. C. Farfan, F. V. Campos.....	381
Diagnosis of occult fractures of the ischiopubic rami	
K. Suzuki.....	391

STIR, SPIR and SPAIR techniques in magnetic resonance of the breast: A comparative study

M. M. Ribeiro, L. Rumor, M. Oliveira, J. G. O’Neill, J. C. Maurício.....395

Advanced magnetic resonance imaging techniques in brain tumours surgical planning

G. Zoccatelli, F. Alessandrini, A. Beltramello, A. Talacchi.....403

Can semi-quantitative evaluation of uncertain (type II) time-intensity curves improve diagnosis in breast DCE-MRI?

R. Fusco, S. Filice, V. Granata, Y. Mandato, A. Porto, M. D’Aiuto, M. Rinaldo, M. Di Bonito, M. Sansone, C. Sansone, A. Rotondo, A. Petrillo.....418

The influence of the blood pressure on the venous cerebral flow measured by magnetic susceptibility (SWI) technique

H. Martins, M. Carreiras, M. M. Ribeiro, P. Sousa, C. Silva-Fortes.....426