

Innovations in Nursing-Driven Bed Resource Optimization for Thoracic Oncology Patients: An Evidence-Based Practice Review

Baowen Huang

Thoracic Department, Sun Yat-sen University Cancer Center, Guangzhou, China

Email: huangbw@sysucc.org.cn

How to cite this paper: Huang, B.W. (2025) Innovations in Nursing-Driven Bed Resource Optimization for Thoracic Oncology Patients: An Evidence-Based Practice Review. *Open Journal of Nursing*, 15, 544-552.
<https://doi.org/10.4236/ojn.2025.157040>

Received: June 19, 2025

Accepted: July 26, 2025

Published: July 29, 2025

Copyright © 2025 by author(s) and Scientific Research Publishing Inc. This work is licensed under the Creative Commons Attribution International License (CC BY 4.0).

<http://creativecommons.org/licenses/by/4.0/>



Open Access

Abstract

Background: Structural inefficiencies in bed allocation for thoracic oncology patients remain a global challenge. This systematic review synthesizes evidence on nurse-led innovations addressing this issue. **Methods:** Following PRISMA 2020 guidelines, we searched PubMed, CNKI, and Wanfang databases (January 2020-March 2025). Quality Appraisal: Two independent reviewers assessed methodological quality using Joanna Briggs Institute (JBI) tools for analytical studies and ROBIS for systematic reviews. Discrepancies were resolved through consensus. 28/32 studies (87.5%) met ≥ 6 JBI criteria, with confounding control being the most frequent limitation. ROBIS indicated low bias risk for 3 systematic reviews. Inclusion criteria covered peer-reviewed studies in English/Chinese reporting quantitative outcomes. Of 1372 initial records, 32 studies met selection criteria after dual screening. Full inclusion criteria: 1) Thoracic oncology focus; 2) Nurse-led bed management intervention; 3) Reported occupancy/transfer/readmission metrics; 4) Controlled before-after or RCT design; 5) ≥ 6 -month implementation. Results: Four evidence-based innovations demonstrated significant impact: 1) Cross-Departmental Admission Criteria Scale reduced unplanned ICU transfers by 27% (95% CI: 22.1 - 31.9; $P < 0.001$) through tri-dimensional risk stratification; 2) ATS-Adapted Handover Protocol decreased clinical errors by 35% ($P = 0.007$) after protocol enhancements; 3) APACHE II-Stratified Nurse Matching maintained unplanned extubation rates at 0.18% (95% CI: 0.15 - 0.22) in high-risk patients; 4) IoT-Enabled Extended Care lowered 30-day readmissions by 18% (RR = 0.82; 95% CI: 0.76 - 0.88). **Conclusions:** Nursing-led bed optimization requires multidimensional integration of risk assessment protocols, technology-enabled surveillance, and policy-supported reimbursement mechanisms.

Keywords

Thoracic Oncology, Bed Resource Allocation, Nursing Risk Stratification, IoT in Healthcare, Value-Based Nursing

1. Introduction: Epidemiological Burden and System Challenges

Thoracic malignancies account for 18% of cancer-related hospitalizations globally, with patients experiencing extended lengths of stay (mean 9.2 ± 2.5 days) due to complex surgical interventions and high complication rates. Meta-analyses indicate postoperative respiratory failure occurs in 12 - 15% of cases, while anastomotic leaks affect 5 - 10% of esophagectomy patients [1]. These clinical challenges are compounded by systemic inefficiencies in bed management.

1.1. Structural Imbalances in Bed Utilization

Analysis of 2024 hospital operational data reveals critical disparities:

- Oncology units consistently operate at >95% capacity, creating bottlenecks for new admissions.

- Adjacent general thoracic surgery departments report 20 - 30% bed vacancy rates.

- Respiratory/ICU departments show intermediate utilization (75 - 85%).

This mismatch generates substantial economic losses. As demonstrated at Xi'an Thoracic Hospital, underutilized beds in general thoracic surgery represented an estimated ¥1.84 million in lost annual revenue, equivalent to supporting 368 additional patient admissions at average reimbursement rates, as shown in **Table 1** [2].

Table 1. 2024 bed utilization metrics in tertiary hospitals.

Department	Occupancy Rate	Vacancy Rate	Revenue Loss ($\times 10^4$ CNY)	Opportunity Cost*
Oncology	96.7%	3.3%	-	-
General Thoracic	73.2%	26.8%	184	368 bed-days
Respiratory & ICU	79.4%	20.6%	127	254 bed-days

*Calculated based on average reimbursement of ¥5000 per bed-day [2].

1.2. The Evolution of Nurse-Led Solutions

Traditional approaches focused on expanding physical capacity, but contemporary models leverage nursing expertise in dynamic resource coordination. Three paradigm shifts characterize modern solutions:

- 1) From Static to Adaptive Allocation: Xi'an Thoracic Hospital's "Whole-Hospital Unified Bed" system employs predictive analytics to anticipate discharges 48 hours in advance, creating flexible bed pools [3].

2) From Task-Oriented to Coordinative Nursing: At Shanxi Bethune Hospital, nurses transformed into “resource orchestrators,” conducting real-time risk assessments during interdepartmental transfers [4].

3) From Paternalistic to Collaborative Care: Mandatory informed consent protocols ensure patients participate in bed allocation decisions, with 100% compliance in documented cases [3].

2. Risk Assessment Innovations: Methodological Advances

2.1. Quality Assessment Protocol

Methodological rigor was evaluated using:

-JBI Critical Appraisal Tools (9-item checklist for quasi-experimental studies) [5] focusing on confounding control, measurement validity, and statistical appropriateness.

-ROBIS (Risk of Bias in Systematic Reviews) [6] assessing domain-specific biases.

Two reviewers independently scored studies ($\kappa = 0.87$). Discrepancies were adjudicated by a third reviewer.

Paper Selection Process: Initial screening (n = 1372) applied:

- (1) Thoracic oncology focus;
- (2) Nurse-led bed management intervention;
- (3) Reported occupancy/transfer/readmission metrics.

Full-text review (n = 89) required:

- (a) Controlled before-after or RCT design;
- (b) ≥ 6 -month implementation;
- (c) Statistical significance reporting (p/CI).

Final inclusion: 32 studies meeting all criteria ($\kappa = 0.87$) [7]-[9].

Results: 28 studies (87.5%) met ≥ 6 JBI criteria. Primary limitations included incomplete confounding control (12 studies) and non-blinded outcome assessment (9 studies). All 3 systematic reviews achieved low ROBIS risk after protocol preregistration (PROSPERO CRD42023487615).

2.2. Handover Protocol Optimization: Technical Enhancements

The ATS checklist was modified through Delphi consensus (42 experts). **Table 2** summarizes key modifications [10].

Table 2. Handover protocol modifications and outcomes.

Original Limitation	Adaptation	Outcome Metric Improvement
Inadequate chylothorax screen	Added rapid triglyceride test (>110 mg/dL)	35% reduction in misclassification
Suboptimal pain assessment	FACES scale for cognitively impaired	28% improvement in pain documentation
Delayed imaging access	PACS-CT integration	86.6% reduction in image retrieval time

Economic Impact Analysis:

- Median handover duration decreased by 6.4 minutes (23.5 → 17.1 min; $P < 0.001$).
- Annual nursing time savings: 4186 hours (equivalent to 2.1 FTE nurses).
- Total cost savings: ¥672,000 including reduced overtime and error correction.

3. Technological Integration in Cross-Departmental Care

3.1. IoT Drainage Monitoring

Hong Kong's smart drainage system demonstrated significant improvements. Clinical outcomes are shown in **Table 3** [4] [11] [12].

System Architecture (Figure 1):

- Edge Computing Gateways (Raspberry Pi 4B) processed data at 2 Hz sampling rate via the MQTT protocol.
- Adaptive Alert Thresholds: Initial drainage >200 mL/h → yellow alert; sustained >100 mL/h → red alert (specificity 98.5%).
- Encryption: TLS 1.3 with SM4 algorithm for audit logs (NIST FIPS 140-2 compliant).

Ethical Implementation: Patient data encryption followed China's GB/T 35273-2020 standard [11]. Consent protocols included:

- 1) Opt-out rights for cloud storage;
- 2) Audit log access;
- 3) Mandatory cybersecurity training (100% nurse compliance).

No breaches reported across 172 patients.




Figure 1. System architecture.

Table 3. IoT system performance metrics (n = 172).

Parameter	Smart System	Manual Recording	Improvement	Security Protocol
Drainage error rate	3.8%	15.2%	75% reduction	TLS 1.3 + SM4
Alert response time	1.7 min	22.4 min	92% faster	NIST FIPS 140-2 compliant
30-day readmission rate	11.7%	29.7%	60.6% lower	Audit log encryption

Cost-Benefit Analysis:

- Device cost: \$238/month (\$2856 annually).
- Avoided readmission savings: \$5210 per case.
- ROI: 1:22.3 (every \$1 invested saves \$22.30 in hospitalization costs).

IoT Technical Configuration

Device specifications are shown in **Table 4**.

Table 4. IoT device configuration settings.

Setting	Device Type	Key Feature
Home Care	Wearable sensor (NFC)	Cloud EHR integration
General Ward	Fixed monitor (5G)	Real-time pressure monitoring
Transitional Care	Hybrid mobile unit	GPS tracking + trend analysis

3.2. Rehabilitation Continuity Framework

Zunyi Medical University's "Three-Phase Standardized Pathway" addresses rehabilitation fragmentation through systematic coordination:

Phase	Core Components	Quality Assurance Mechanism
Pre-op	30-minute daily modified Yijin Jing	Therapist-supervised sessions
Transfer	Rehab progress documentation	Dual-nurse verification
Post-transfer	OSCE-validated competency assessment	≥90% accuracy requirement

Physiological Outcomes:

- Postoperative Day 3 FEV₁: 58.3% predicted vs. 42.1% in controls (Δ 16.2%, $P = 0.002$) [8].
- Rehabilitation interruption rate: 11.7% vs. 31.6% pre-implementation (63% reduction, $P < 0.001$) [8].
- Patient-reported adherence: 89.2% for simplified exercise protocol.

4. Human Resource Optimization Models**4.1. APACHE II-Stratified Staffing: Implementation Framework**

The competency-matching model employs evidence-based risk stratification. The operational protocol is detailed in **Table 5** and **Figure 2**.

Performance Outcomes:

- Unplanned extubation rate: 0.18% (95% CI: 0.14 - 0.23) vs. target $\leq 0.2\%$.
- Remote consultation success: 94.3% (95% CI: 91.7 - 96.2) for drain complications.
- Training efficiency: 89% pass rate for ECMO management program ($n = 112$ nurses).




Figure 2. Operational protocol.

4.2. Burnout Mitigation: Multidimensional Approach

Nanjing Medical University's intervention targeted burnout through three evidence-based strategies:

Table 5. Comparative effectiveness.

Strategy	Implementation Detail	Outcome Change	Effect Size
Cross-training	Quarterly 2-week rotations	OSCE pass rate ↑ 76% → 98%	Cohen's d = 1.2
Competency matching	APACHE II-driven assignments	Job-person fit ↑ 3.2 → 4.5 (of 5.0)	$r^2 = 0.37$

Burnout Reduction:

-MBI emotional exhaustion: Pre 28.7 ± 6.2 → Post 20.6 ± 5.4 ($\Delta 28.2\%$, $P < 0.001$).

-12-month retention: 93.4% vs. 81.7% pre-intervention.

4.3. Nursing Challenges in Bed Coordination

Implementation Barriers:

-Resistance to change: 42% of nurses (95% CI: 38 - 46) opposed APACHE II stratification [8].

-Training gaps: 68% lacked competency in cross-departmental parameter interpretation (e.g., neurological scales) [13].

-Resource constraints: Rural 5G coverage <40% vs. urban 95% (MIIT 2024) [14].

Solutions:

-Microtraining: Mobile modules reduced knowledge gaps by 41% (Δ pre/post-test scores: $58.3 \rightarrow 82.1$, $P < 0.001$).

-AI Skill Matching: Zhejiang Nursing Cloud platform improved job-person fit from 3.2 to 4.1/5.0 ($r^2 = 0.29$).

5. Extended Care and Policy Integration

5.1. Limitations and Generalizability

Geographical bias exists with 26/32 (81.3%) studies from Chinese single centers. Generalizability may be constrained by:

- Higher nurse-patient ratios (1:3 vs. OECD average 1:6.2).
- Advanced IoT infrastructure (urban 5G coverage 95% vs. EU 68%).

Quasi-experimental designs enhanced feasibility but limited causal inference. Validation in diverse systems (e.g., US safety-net hospitals) is warranted.

5.2. Policy Integration Enabling Scalability

Alignment with Article 12 of China's Internet Plus Nursing Services enabled Zhejiang's payment model (ZJ-TB-2024) [15], reducing out-of-pocket costs by 57%.

Long-Term Viability: Zhejiang's ZJ-TB-2024 model demonstrated sustainability through:

- Reimbursement redesign: Bundled payments for IoT monitoring (¥420/patient).
- Infrastructure grants: Provincial 5G expansion to 80% rural coverage by 2026.
- Outcome-linked funding: 18% readmission reduction → 7% bonus payments.

5.3. Evidence Synthesis

Recent systematic reviews confirm nurse-led bed allocation reduces ICU overflow (OR = 0.71, 95% CI: 0.63 - 0.81) but emphasize interoperability standards as critical success factors [1].

6. Conclusions: Strategic Implications and Research Agenda

The transformation of thoracic oncology nursing represents a paradigm shift from compartmentalized care to integrated resource coordination. Three pillars support this evolution:

6.1. Evidence-Based Transformation Pathways

-Decision-Support Tools: The tri-dimensional admission scale exemplifies how standardized assessment can reduce practice variation while improving outcomes (unplanned ICU transfer ↓ 27%) [7].

-Technology Enablement: IoT systems create virtual care continuums, reducing readmissions through early detection (RR = 0.82) [12].

-Policy Alignment: Innovative payment models like ZJ-TB-2024 [15] demonstrate how reimbursement reform enables sustainable scaling.

6.2. Implementation Science Priorities

Future research should address critical evidence gaps:

-Health Economics: Cost-effectiveness analysis of IoT monitoring across healthcare systems (e.g., ICER per QALY gained).

-Model Adaptability: Validation of APACHE II staffing models in community hospitals.

-Digital Infrastructure: Development of cloud-based nursing coordination platforms with interoperability standards.

6.3. Nursing Leadership Imperatives

-Advocate for inclusion in hospital resource allocation committees.

-Develop metrics demonstrating nursing's impact on operational efficiency.

-Establish cross-institutional learning collaboratives for best practice diffusion.

Conflicts of Interest

The author declares no conflicts of interest regarding the publication of this paper.

References

- [1] Smith, R., Johnson, A., Williams, D., Brown, S., Miller, J., Davis, P., *et al.* (2024) Nurse-Led Bed Management in Oncology: A Systematic Review. *Journal of Healthcare Management*, **39**, e1-e15.
- [2] Xi'an Thoracic Hospital Operations Department (2025) 2024 Annual Report on Bed Resource Utilization and Operational Efficiency [Internal Report]. XTH Press.
- [3] Shanxi Bethune Hospital Cancer Center (2024) Multicenter Study on Extended Nursing for Thoracic Oncology Patients [Technical Report]. Shanxi Bethune Hospital.
- [4] Liu, X. and Chen, H. (2025) Development and Validation of APACHE II-Stratified Nursing Pathways for Lung Cancer Patients. *Journal of Nursing Science*, **40**, 1-5.
- [5] Zhang, L., Wang, Y., Zhao, M., Sun, W., Chen, X., Li, J., *et al.* (2024) Derivation and Validation of a Prediction Model for Unplanned ICU Transfer after Thoracic Surgery. *Chest*, **166**, 1120-1128.
- [6] Wong, T., Chan, K.F., Lee, S., Lam, T.N., Ho, W.S., Cheng, M., *et al.* (2025) Real-Time IoT Monitoring System for Postoperative Thoracic Drainage: A Randomized Controlled Trial. *The Annals of Thoracic Surgery*, **109**, 412-418.
- [7] National Nursing Quality Data Platform (2025) Optimizing Handover Processes to Enhance Patient Safety in Thoracic Surgery. *Chinese Nursing Management*, **25**, 45-49.
- [8] Thoracic Nursing Committee, Chinese Nursing Association (2024) Standardized Handover Protocol for Thoracic Oncology Patients: Expert Consensus. *Chinese Journal of Nursing*, **59**, 289-295.
- [9] Huang, F., Zhou, M., Zheng, R., Wu, X.L., Lin, L., Cai, H., *et al.* (2025) Cross-Departmental Competency Training for Oncology Nurses. *Journal of Clinical Nursing*, **34**, 123-130.
- [10] Thoracic Nursing Committee, Chinese Nursing Association (2024) Enhanced ATS Handover Protocol: Technical Specifications. CMA Press.
- [11] General Administration of Market Supervision (2020) GB/T 35273-2020 Information Security Technology—Personal Information Security Specification. Standards Press.
- [12] Chinese Association of Thoracic Surgeons (2023) Clinical Guidelines for Prevention and Management of Thoracic Surgery Complications. People's Medical Publishing House.

- [13] MIIT China (2025) 5G Network Coverage Report 2024. Ministry of Industry and IT.
- [14] National Health Commission (2025) 2025 National Healthcare IoT Implementation Guidelines. NHC Press.
- [15] Zhejiang Medical Insurance Bureau (2024) ZJ-TB-2024 Payment Model for IoT-Enabled Nursing Services. ZJ Health Press.